



文 献

第1章～特典

参考文献

- (1) Ueshima H. Explanation for the Japanese paradox: prevention of increase in coronary heart disease and reduction in stroke. *J Atheroscler Thromb.* 2007; 14: 278-86.
- (2) Kubo M, 他. Trends in the incidence, mortality, and survival rate of cardiovascular disease in a Japanese Community:the Hisayama study. *Stroke.* 2003; 34: 2349-54.
- (3) 介護保険事業状況報告 [Internet]. [cited 2018 Feb 26]. Available from: <https://www.e-stat.go.jp/stat-search/files?page=1&toukei=00450351&tstat=000001031648>
- (4) 国民生活基礎調査 [Internet]. [cited 2018 Feb 2]. Available from: <https://www.e-stat.go.jp/stat-search/files?page=1&toukei=00450061&tstat=000001031016>
- (5) Yoshida D, Ninomiya T, Doi Y, Hata J, Fukuhara M, Ikeda F, Mukai N, Kiyohara Y. Prevalence and causes of functional disability in an elderly general population of Japanese: the Hisayama study. *J Epidemiol.* 2012; 22: 222-9.
- (6) Takashima N, Arima H, Kita Y, Fujii T, Miyamatsu N, Komori M, Sugimoto Y, Nagata S, Miura K, Nozaki K. Incidence, management and short-term outcome of stroke in a general population of 1.4 million Japanese- Shiga Stroke Registry. *Circ J.* 2017.
- (7) Kucharska-Newton A, Griswold M, Yao ZH, Foraker R, Rose K, Rosamond W, Wagenknecht L, Koton S, Pompeii L, Windham BG. Cardiovascular Disease and Patterns of Change in Functional Status Over 15 Years: Findings From the Atherosclerosis Risk in Communities (ARIC) Study. *J Am Heart Assoc.* 2017; 6.
- (8) Dhamoon MS, Longstreth WT, Jr., Bartz TM, Kaplan RC, Elkind MSV. Disability trajectories before and after stroke and myocardial infarction: The Cardiovascular Health Study. *JAMA Neurol.* 2017; 74: 1439-45.
- (9) Hozawa A, Okamura T, Murakami Y, Kadokawa T, Okuda N, Takashima N, Hayakawa T, Kita Y, Miura K, Nakamura Y, Okayama A, Ueshima H, Group NDR. High blood pressure in middle age is associated with a future decline in activities of daily living. *NIPPON DATA 80. J Hum Hypertens.* 2009; 23: 546-52.
- (10) Takashima N, Miura K, Hozawa A, Okamura T, Hayakawa T, Okuda N, Kadokawa T, Murakami Y, Kita Y, Nakamura Y, Okayama A, Ueshima H, Group NDR. Cigarette smoking in middle age and a long-term risk of impaired activities of daily living: NIPPON DATA 80. *Nicotine Tob Res.* 2010; 12: 944-9.
- (11) Hayakawa T, Okamura T, Okayama A, Kanda H, Watanabe M, Kita Y, Miura K, Ueshima H. Relationship between 5-year decline in instrumental activity of daily living and accumulation of cardiovascular risk factors: NIPPON DATA 90. *J Atheroscler Thromb.* 2010; 17: 64-72.
- (12) 日本老年医学会. フレイルに関する日本老年医学会からのステートメント 2014 [updated 2014 May; cited 2018 Feb 10]. Available from: https://www.jpn-geriat-soc.or.jp/info/topics/pdf/20140513_01_01.pdf
- (13) Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, Seeman T, Tracy R, Kop WJ, Burke G, McBurnie MA, Cardiovascular Health Study Collaborative Research G. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci.* 2001; 56: M146-56.
- (14) Kojima G, Iliffe S, Taniguchi Y, Shimada H, Rakugi H, Walters K. Prevalence of frailty in Japan: A systematic review and meta-analysis. *J Epidemiol.* 2017; 27: 347-53.
- (15) Satake S, Shimada H, Yamada M, Kim H, Yoshida H, Gondo Y, Matsubayashi K, Matsushita E, Kuzuya M, Kozaki K, Sugimoto K, Senda K, Sakuma M, Endo N, Arai H. Prevalence of frailty among community-dwellers and outpatients in Japan as defined by

- the Japanese version of the Cardiovascular Health Study criteria. *Geriatr Gerontol Int.* 2017; 17: 2629-34.
- (16) Makizako H, Shimada H, Doi T, Tsutsumimoto K, Suzuki T. Impact of physical frailty on disability in community-dwelling older adults: a prospective cohort study. *BMJ Open.* 2015; 5: e008462.
 - (17) Ohara T, Hata J, Yoshida D, Mukai N, Nagata M, Iwaki T, Kitazono T, Kanba S, Kiyohara Y, Ninomiya T. Trends in dementia prevalence, incidence, and survival rate in a Japanese community. *Neurology.* 2017; 88: 1925-32.
 - (18) Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D, Ballard C, Banerjee S, Burns A, Cohen-Mansfield J, Cooper C, Fox N, Gitlin LN, Howard R, Kales HC, Larson EB, Ritchie K, Rockwood K, Sampson EL, Samus Q, Schneider LS, Selbaek G, Teri L, Mukadam N. Dementia prevention, intervention, and care. *Lancet.* 2017; 390: 2673-734.
 - (19) Fujiyoshi A, Ohkubo T, Miura K, Murakami Y, Nagasawa SY, Okamura T, Ueshima H. Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. *Hypertens Res.* 2012; 35: 947-53.
 - (20) Takashima N, Ohkubo T, Miura K, Okamura T, Murakami Y, Fujiyoshi A, Nagasawa SY, Kadota A, Kita Y, Miyagawa N, Hisamatsu T, Hayakawa T, Okayama A, Ueshima H. Long-term risk of BP values above normal for cardiovascular mortality: a 24-year observation of Japanese aged 30 to 92 years. *J Hypertens.* 2012; 30: 2299-306.
 - (21) Ikeda A, Iso H, Yamagishi K, Inoue M, Tsugane S. Blood pressure and the risk of stroke, cardiovascular disease, and all-cause mortality among Japanese: the JPHC Study. *Am J Hypertens.* 2009; 22: 273-80.
 - (22) Arima H, Tanizaki Y, Yonemoto K, Doi Y, Ninomiya T, Hata J, Fukuhara M, Matsumura K, Iida M, Kiyohara Y. Impact of blood pressure levels on different types of stroke: the Hisayama study. *J Hypertens.* 2009; 27: 2437-43.
 - (23) Imano H, Kitamura A, Sato S, Kiyama M, Ohira T, Yamagishi K, Noda H, Tanigawa T, Iso H, Shimamoto T. Trends for blood pressure and its contribution to stroke incidence in the middle-aged Japanese population: the Circulatory Risk in Communities Study (CIRCS). *Stroke.* 2009; 40: 1571-7.
 - (24) Murakami Y, Hozawa A, Okamura T, Ueshima H. Relation of blood pressure and all-cause mortality in 180,000 Japanese participants: pooled analysis of 13 cohort studies. *Hypertension.* 2008; 51: 1483-91.
 - (25) Ikeda N, Saito E, Kondo N, Inoue M, Ikeda S, Satoh T, Wada K, Stickley A, Katanoda K, Mizoue T, Noda M, Iso H, Fujino Y, Sobue T, Tsugane S, Naghavi M, Ezzati M, Shibuya K. What has made the population of Japan healthy? *Lancet.* 2011; 378: 1094-105.
 - (26) Turin TC, Murakami Y, Miura K, Rumana N, Kita Y, Hayakawa T, Okamura T, Okayama A, Ueshima H. Hypertension and life expectancy among Japanese: NIPPON DATA80. *Hypertens Res.* 2012; 35: 954-8.
 - (27) 厚生労働省. 平成28年国民健康・栄養調査報告; 2016.
 - (28) 三浦克之(研究代表者). 厚生労働行政推進調査事業費補助金・循環器疾患・糖尿病等生活習慣病対策総合研究事業「新旧(1980-2020年)ライフスタイルからみた国民代表集団大規模コホート研究: NIPPON DATA80/90/2010/2020(H30-循環器等-指定-002)」平成30年度総括・分担研究報告書 2019.
 - (29) Stamler J, Elliott P, Chan Q. INTERMAP appendix tables. *J Hum Hypertens.* 17: 665-775. 2003.

- (30) Nakamura K, Okamura T, Hayakawa T, Hozawa A, Kadokawa T, Murakami Y, et al. The proportion of individuals with obesity-induced hypertension among total hypertensives in a general Japanese population: NIPPON DATA80, 90. *Eur J Epidemiol* 2007; 22: 691-698.
- (31) Neter JE, Stam BE, Kok EJ, et al. Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 42: 878-884, 2003.
- (32) Marmot MG, Elliott P, Shipley MJ, Dyer AR, Ueshima H, Beevers DG, et al. Alcohol and blood pressure: the INTERSALT study. *BMJ* 1994; 308: 1263-1267.
- (33) Yoshita K, Miura K, Morikawa Y, Ishizaki M, Kido T, Naruse Y, et al. Relationship of alcohol consumption to 7 year blood pressure change in Japanese men. *J Hypertens* 2005;23:1485-1490.
- (34) Okamura T, Tanaka T, Yoshita K, Chiga N, Takebayashi T, Kikuchi Y, et al. Specific alcoholic beverage and blood pressure in a middle-aged Japanese population: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) Study. *J Hum Hypertens* 2004; 18: 9-16.
- (35) Witteman JC, Willett WC, Stampfer MJ, Colditz GA, Kok FJ, Sacks FM, et al. Relation of moderate alcohol consumption and risk of systemic hypertension in women. *Am J Cardiol* 1990; 65: 633-637.
- (36) Tsuruta M, Adachi H, Hirai Y, Fujiura Y, Imaizumi T. Association between alcohol intake and development of hypertension in Japanese normotensive men: 12-year follow-up study. *Am J Hypertens* 2000; 13(5 Pt 1):482-487.
- (37) Ueshima H, Mikawa K, Baba S, Sasaki S, Ozawa H, Tsushima M, et al. Effect of reduced alcohol consumption on blood pressure in untreated hypertensive men. *Hypertension* 1993; 21: 248-252.
- (38) Xin X, He J, Frontini MG, Ogden LG, Motsamai OI, Whelton PK. Effects of alcohol reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 2001;38: 1112-1117.
- (39) Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbiagele B, Smith SC Jr, Spencer CC, Stafford RS, Taler SJ, Thomas RJ, Williams KA Sr, Williamson JD, Wright JT Jr. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/ APhA/ASH/ ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018; 71(6): e13-e115.
- (40) 日本高血圧学会高血圧治療ガイドライン作成委員会編. 高血圧治療ガイドライン2019. 日本高血圧学会, 2019.
- (41) Kiyanaga A, Arakawa K, Tanaka H, Shindo M. Blood pressure and hormonal responses to aerobic exercise. *Hypertension* 1985; 7: 125-131.
- (42) Welton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized controlled trials. *Ann Intern Med* 2002; 136: 493-503.
- (43) Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *J Am Heart Assoc.* 2013; 2: e004473.
- (44) Dahl LK. Possible role of salt intake in the development of essential hypertension, in Cottier P, Bock KD (eds): *Essential Hypertension - An International Symposium*. Berlin,

- Springer-Verlag, 1960. pp 53-65.
- (45) Intersalt Cooperative Research Group. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. *BMJ* 1988; 297: 319-328.
- (46) Cutler JA, Follmann D, Allender PS. Randomized trials of sodium reduction: an overview. *Am J Clin Nutr* 1997; 65(suppl): 643S-651S.
- (47) He FJ, MacGregor GA. Effect of modest salt reduction on blood pressure: a meta-analysis of randomized trials. Implications for public health. *J Hum Hypertens* 2002; 16: 761-770.
- (48) Appel LJ, Moore TJ, Obarzanek E, et al. A clinical trial of the effects of dietary patterns on blood pressure. *N Engl J Med* 1997; 336:1117-1124.
- (49) Miura K, Torii S. Diet, Nutrients, and the Prevention of Hypertension. *Curr Nutr Rep* 2012; DOI 10.1007/s13668-012-0012-4.
- (50) UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS33). *Lancet* 1998; 352: 837-53.
- (51) The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med.* 1993; 329: 977-86.
- (52) Zhang X et al.; Prevalence of diabetic retinopathy in the United States, 2005-2008. *JAMA* 304: 649-56, 2010.
- (53) Kawasaki R, et al. ; Japan Diabetes Complications Study Group: Incidence and progression of diabetic retinopathy in Japanese adults with type 2 diabetes: 8 year follow-up study of the Japan Diabetes Complications Study (JDGS). *Diabetologia*. 54: 2288-94, 2011.
- (54) 中江公裕, 他:わが国における視覚障害の現状. 厚生労働省難治性疾患克服研究事業 網脈絡膜・視神經萎縮症に関する研究:平成17年度研究報告書. 263-267, 2006.
- (55) 一般社団法人日本透析医学会(<http://www.jsdt.or.jp/>) : 図説 わが国の慢性透析療法の現況
- (56) 日本糖尿病学会:糖尿病網膜症の治療. 科学的根拠に基づく糖尿病診療ガイドライン2013.
- (57) Yokoyama H, et al. Microalbuminuria is common in Japanese: Type 2 Diabetic Patients. A nationwide survey from the Japan Diabetes Clinical Data Management Study Group (JDDM 10). *Diabetes Care*, 30: 989-92, 2007.
- (58) Katayama S, et al. for the Japan Diabetes Complications Study Group. Low transition rate from normo- and low Microalbuminuria to proteinuria in Japanese type 2 diabetic individuals: the Japan Diabetes Complications Study (JDGS). *Diabetologia* 54: 1025-31, 2011.
- (59) Molitch ME, et al.; Diabetic nephropathy. American Diabetes Association. *Diabetes Care*. 26 Suppl 1: S94-8, 2003.
- (60) Ritz E, et al.; Nephropathy in patients with type 2 diabetes mellitus. *N Engl J Med.* 341: 1127-33, 1999.
- (61) 日本糖尿病対策推進会議(<http://www.med.or.jp/tounyoubyou/>):日本における糖尿病患者の足外観異常, および, 糖尿病性神経障害の実態に関する報告. 平成20年3月.
- (62) The DECODE study group on behalf of the European Diabetes Epidemiology Group: Glucose tolerance and mortality: comparison of WHO and American. *Lancet*; 354: 617-21, 1999.

- (63) The DECODE study group on behalf of the European Diabetes Epidemiology Group : Is the Current Definition for Diabetes Relevant to Mortality Risk From All Causes and Cardiovascular and Noncardiovascular Diseases? *Diabetes Care* 26: 688-696, 2003.
- (64) JL Chiasson, et al. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet*. 359, 2072-2077, 2002.
- (65) JL Chiasson, et al. Acarbose treatment and the risk of cardiovascular disease and hypertension in patients with impaired glucose tolerance: the STOP-NIDDM trial. *JAMA*. 290, 486-494, 2003.
- (66) Doi Y, et al. Impact of Glucose Tolerance Status on Development of Ischemic Stroke and Coronary Heart Disease in a General Japanese Population: The Hisayama Study. *Stroke* 2010; 41: 203-9.
- (67) Kadowaki S, et al. Relationship of elevated casual blood glucose level with coronary heart disease, cardiovascular disease and all-cause mortality in a representative sample of the Japanese population. *NIPPON DATA80*. *Diabetologia* 2008; 51: 575-82.
- (68) Kokubo Y, et al. The combined impact of blood pressure category and glucose abnormality on the incidence of cardiovascular diseases in a Japanese urban cohort: the Suita Study. *Hypertens Res* 2010; 33: 1238-43.
- (69) Watanabe M, et al. New diagnosis criteria for diabetes with hemoglobin A1c and risks of macro-vascular complications in an urban Japanese cohort: The Suita Study. *Diabetes Res Clin Pract* 2010; 88: e20-3.
- (70) Watanabe M, et al. Serum 1,5-anhydro-d-glucitol levels predict first-ever cardiovascular disease: An 11-year population-based Cohort study in Japan, the Suita Study. *Atherosclerosis* 2011; 216: 477-83.
- (71) Holman RR et al. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008; 359: 1577-1589.
- (72) Nathan DM, et al.; Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Study Research Group. Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med*; 353: 2643-53, 2005.
- (73) Whiting DR, et al.; IDF Diabetes Atlas: Global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Res Clin Pract* 2011; 94: 311-21.
- (74) International Diabetes Federation(IDF : 国際糖尿病連合) ウェブサイト (<http://www.idf.org/>)
- (75) 厚生省保険医療局(<http://www.mhlw.go.jp/>) : 平成9年糖尿病実態調査(平成11年4月).
- (76) 厚生労働省健康局(<http://www.mhlw.go.jp/>) : 平成14年度糖尿病実態調査報告(平成16年6月).
- (77) 厚生労働省健康局(<http://www.mhlw.go.jp/>) : 平成19年国民健康・栄養調査報告.
- (78) 厚生労働省健康局(<http://www.mhlw.go.jp/>) : 平成24年国民健康・栄養調査報告.
- (79) 厚生労働省健康局(<http://www.mhlw.go.jp/>) : 平成28年国民健康・栄養調査報告.
- (80) Pan XR, et al. Effects of Diet and Exercise in Preventing NIDDM in People With Impaired Glucose Tolerance. The Da Qing IGT and Diabetes Study. *Diabetes care* 20: 537-44, 1997.
- (81) Tuomilehto J, et al. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance, *N Engl J Med*; 344: 1343-50. 2001.
- (82) Knowler WC, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin, *N Engl J Med*; 346: 393-403. 2002.

- (83) Ramachadran A et al. The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia.*; 49: 289-97. 2006.
- (84) Saito T, et al. Lifestyle modification and prevention of type 2 diabetes in overweight Japanese with Impaired fasting glucose levels A randomized controlled trial. *Arch Intern Med.* 2011; 171 (15): 1352-1360.
- (85) 健康・栄養情報研究会：国民栄養の現状(平成14年厚生労働省国民栄養調査結果)，第一出版。
- (86) JM. Chan, et al.: Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes care* 17: 961-9, 1994.
- (87) Colditz GA, et al.: Weight gain as a risk factor for clinical diabetes mellitus in women. *Ann Intern Med.*; 122: 481-6, 1995.
- (88) Carey VJ, et al.: Body Fat Distribution and Risk of Non-Insulin-dependent Diabetes Mellitus in Women. The Nurses' Health Study. *Am J Epidemiol.* 145: 614-9, 1997.
- (89) Sone H, et al. Energy intake and obesity in Japanese patients with type 2 diabetes. *Lancet.* 17; 363 (9404): 248-9, 2004.
- (90) Jeon CY, et al. Physical activity of moderate intensity and risk of type 2 diabetes: a systematic review *Diabetes Care* 30: 744-52, 2007.
- (91) Okada K, et al.: Leisure-time physical activity at weekends and the risk of Type 2 diabetes mellitus in Japanese men: the Osaka Health Survey. *Diabet Med* 17: 53-58, 2000.
- (92) Nakanishi N, et al.: Daily life activity and risk of developing impaired fasting glucose or Type2 diabetes in middel-aged Japanese. *Diabetologia* 47: 1768-1775, 2004.
- (93) Carlsson S, et al. Alcohol consumption and type 2 diabetes: metaanalysis of epidemiological studies indicates a U-shaped relationship. *Diabetologia* 48: 1051-1054, 2005.
- (94) Koppes LL, et al. Bouter LM, Heine RJ. Moderate alcohol consumption lowers the risk of type 2 diabetes: a meta-analysis of prospective observational studies. *Diabetes Care* 28: 719-25, 2005.
- (95) Baliunas DO, et al. Alcohol as a risk factor for type 2 diabetes: A systematic review and meta-analysis. *Diabetes Care* 32: 2123-32, 2009.
- (96) Tsumura K, et al.: Daily alcohol consumption and the risk of Type 2 diabetes in Japanese Men. The Osaka Health Survey. *Diabetes care* 22: 1432-1437, 1999.
- (97) Watanabe M, et al. Alcohol consumption and the risk of diabetes by body mass index levels in a cohort of 5636 Japanese. *Diabetes Res Clin Pract.*; 57: 191-7, 2002.
- (98) Waki K, et al. for the JPHC Study Group. Alcohol consumption and other risk factors for self-reported diabetes among middle-aged Japanese: a population-based prospective study in the JPHC study cohort I. *Diabet Med* 22: 323-31, 2005.
- (99) Nakanishi N, et al. Alcohol consumption and risk for development of impaired fasting glucose or type 2 diabetes in middle-aged Japanese men. *Diabetes Care.* 2003; 26: 48-54.
- (100) Willi C. Active Smoking and the Risk of Type 2 Diabetes. A Systematic Review and Meta-analysis. *JAMA.* 2007; 298 (22): 2654-64.
- (101) Oba S, et al. Smoking cessation increases short-term risk of type 2 diabetes irrespective of weight gain: the Japan Public Health Center-Based Prospective Study. *PLoS One.* 2012; 7 (2): e17061.
- (102) Okamura T, Tanaka H, Miyamatsu N, et al. The relationship between serum total cholesterol and all-cause or cause-specific mortality in a 17.3-year study of a Japanese

- cohort. *Atherosclerosis* 190: 216-23, 2007.
- (103) Cui R, Iso H, Toyoshima H, et al. Serum total cholesterol levels and risk of mortality from stroke and coronary heart disease in Japanese: the JACC study. *Atherosclerosis* 194: 415-20, 2007.
 - (104) Okamura T, Kokubo Y, Watanabe M, et al. Low-density-lipoprotein cholesterol and non-high density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort study: the Suita study. *Atherosclerosis* 203: 587-92, 2009.
 - (105) Imano H, et al. Low-density lipoprotein cholesterol and risk of coronary heart disease among Japanese men and women: The Circulatory Risk in Communities Study (CIRCS). *Prev Med* 52: 381-6, 2011.
 - (106) Nagasawa SY, Okamura T, Iso H, et al. Relation Between Serum Total Cholesterol Level and Cardiovascular Disease Stratified by Sex and Age Group: A Pooled Analysis of 65 594 Individuals From 10 Cohort Studies in Japan. *J Am Heart Assoc*; 1: e001974, 2012.
 - (107) Kitamura A, Iso H, Naito Y, et al. High-density lipoprotein cholesterol and premature coronary heart disease in urban Japanese men. *Circulation*;89: 2533-9, 1994.
 - (108) Briel M, Ferreira-Gonzalez I, You JJ , et al. Association between change in high density lipoprotein cholesterol and cardiovascular disease morbidity and mortality: systematic review and meta-regression analysis. *BMJ* 338: b92, 2009.
 - (109) The AIM-HIGH Investigators. Niacin in Patients with Low HDL Cholesterol Levels Receiving Intensive Statin Therapy. *New Engl J Med* 2011; 365: 2255-2267.
 - (110) Barter PJ, Caulfield M, Eriksson M, Grundy SM, Kastelein JJ, Komajda M, Lopez-Sendon J, Mosca L, Tardif JC, Waters DD, Shear CL, Revkin JH, Buhr KA, Fisher MR, Tall AR, Brewer B; ILLUMINATE Investigators. Effects of torcetrapib in patients at high risk for coronary events. *N Engl J Med* 2007; 357: 2109-22.
 - (111) Hirata T, Sugiyama D, Nagasawa SY, Murakami Y, Saitoh S, Okayama A, Iso H, Irie F, Sairenchi T, Miyamoto Y, Yamada M, Ishikawa S, Miura K, Ueshima H, Okamura T. A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan. *Eur J Epidemiol* 2017; 32: 547-557.
 - (112) Iso H, Naito Y, Sato S, et al. Serum triglycerides and risk of coronary heart disease among Japanese men and women. *Am J Epidemiol* 153: 490-499, 2001.
 - (113) Okamura T, Kokubo Y, Watanabe M, et al. Triglycerides and non-high-density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort: the Suita study. *Atherosclerosis* 209: 290-4, 2010.
 - (114) Tanaka T, Okamura T. Blood cholesterol level and risk of stroke in community-based or worksite cohort studies: a review of Japanese cohort studies in the past 20 years. *Keio J Med* 61: 79-88, 2012.
 - (115) Imamura T, Doi Y, Arima H, et al. LDL cholesterol and the development of stroke subtypes and coronary heart disease in a general Japanese population: the Hisayama study. *Stroke*; 40: 382-8, 2009.
 - (116) Nakamura H, Arakawa K, Itakura H, et al. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *Lancet*; 368 (9542): 1155-63, 2006.
 - (117) Kushiro T, Mizuno K, Nakaya N, et al. Pravastatin for cardiovascular event primary prevention in patients with mild-to-moderate hypertension in the Management of

- Elevated Cholesterol in the Primary Prevention Group of Adult Japanese (MEGA) Study. Hypertension 53: 135-41, 2009.
- (118) Amarenco P, Bogousslavsky J, Callahan A 3rd, et al. High-dose atorvastatin after stroke or transient ischemic attack. N Engl J Med 355: 549-59, 2006.
- (119) Hosomi N, Nagai Y, Kohriyama T, Ohtsuki T, Aoki S, Nezu T, Maruyama H, Sunami N, Yokota C, Kitagawa K, Terayama Y, Takagi M, Ibayashi S, Nakamura M, Origasa H, Fukushima M, Mori E, Minematsu K, Uchiyama S, Shinohara Y, Yamaguchi T, Matsumoto M; J-STARS Collaborators. The Japan Statin Treatment Against Recurrent Stroke (J-STARS): A Multicenter, Randomized, Open-label, Parallel-group Study. EBioMedicine 2015; 2: 1071-8.
- (120) Jacobs D, Blackburn H, Higgins M, et al. Report of the conference on low blood cholesterol: mortality associations. Circulation, 86: 1046-1060, 1992.
- (121) Hirano K, Yamashita S, Nakajima N, et al. Genetic Cholesteryl Ester Transfer Protein deficiency is extremely frequent in Omagari area of Japan. Marked hyperalphalipoproteinemia caused by CETP gene mutation is not associated with longevity. Arterioscler Thromb Vasc Biol 17: 1053-1059, 1997.
- (122) Moriyama Y, Okamura T, Inazu A, et al.: A Low Prevalence of Coronary Heart Disease among Subjects with Increased High-Density Lipoprotein Cholesterol Levels, Including Those with Plasma Cholesteryl Ester Transfer Protein Deficiency. Prev Med 27: 659-667, 1998.
- (123) Curb JD, Abbott RD, Rodriguez BL, et al. A prospective study of HDL cholesterol and cholesteryl ester transfer protein gene mutations and the risk of coronary heart disease in the elderly. J Lipid Res 2004; 45: 948-53.
- (124) Hirata A, Sugiyama D, Watanabe M, Tamakoshi A, Iso H, Kotani K, Kiyama M, Yamada M, Ishikawa S, Murakami Y, Miura K, Ueshima H, Okamura T. Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of nine cohort studies including 43,407 individuals: the EPOCH-JAPAN study. J Clin Lipidol, 12: 674-684, 2018.
- (125) Verschuren WM, Jacobs DR, Bloomberg BP, et al. Serum total cholesterol and long-term coronary heart disease mortality in different cultures. Twenty-five-year follow-up of the seven countries study. JAMA. 274: 131-136, 1995.
- (126) 関川暁, 上島弘嗣, Curb DJ, Kuller LH. 21世紀の国際間比較対照研究 ERA-JUMP予防循環器病における日本の挑戦, 世界への貢献. 医学のあゆみ207: 482-486, 2003.
- (127) 厚生労働省. 平成28年国民健康・栄養調査報告(<https://www.mhlw.go.jp/bunya/kenkou/eiou/dl/h28-houkoku.html>)
- (128) 国立健康・栄養研究所：国民健康・栄養の現状－平成29年厚生労働省国民健康・栄養調査報告より－. 第一出版株式会社, 2019.
- (129) Guo Z, Miura K, Turin TC, et al. Relationship of the polyunsaturated to saturated fatty acid ratio to cardiovascular risk factors and metabolic syndrome in Japanese: the INTERLIPID study. J Atheroscler Thromb. 17(8):777-84, 2010.
- (130) 岡村智教, 東山綾, 宮松直美. 喫煙による低HDL血症. 脂質代謝異常－高脂血症・低脂血症－. 日本臨床(増刊号)65: 663-667, 2007.
- (131) Imamura H, et al.: Relationship of cigarette smoking with blood pressure, serum lipids and lipoproteins in young Japanese women. Clin Exp Pharmacol Physiol 27: 364-369, 2000.
- (132) Kuzuya M, et al.: Effect of smoking habit on age-related changes in serum lipids: A cross-

- sectional and longitudinal analysis in a large Japanese cohort. *Atherosclerosis* 185: 183-190, 2006.
- (133) Bernaards CM, et al.: In a prospective study in young people, associations between changes in smoking behavior and risk factors for cardiovascular disease were complex. *J Clin Epidemiol* 58: 1165-1171, 2005.
- (134) Maeda K, et al.: The effects of cessation from cigarette smoking on the lipid and lipoprotein profiles: a meta-analysis. *Prev Med* 37: 283-290, 2003.
- (135) Moffatt RJ, et al.: Effects of the Transdermal Nicotine Patch on Normalization of HDL-C and Its Subfractions. *Prev Med* 31: 148-152, 2000.
- (136) 岡村智教, 早川岳人, 門脇 崇, 天本健司, 喜多義邦, 岡山 明, 上島弘嗣. 日本人の代表集団における飲酒量の変化に伴う循環器疾患死亡の相対リスク, 絶対リスク変化量の推計-NIPPON DATA-. 日本アルコール・薬物医学会雑誌 36: 586-595, 2001.
- (137) 岡村智教ほか: HDLコレステロールと歩行習慣の関連-歩数計を用いた検討. *動脈硬化* 21: 585-589, 1994.
- (138) Choudhury SR, et al. Alcohol intake and serum lipids in a Japanese population. *Int J Epidemiol* 23: 940-7, 1994.
- (139) Higashiyama A, et al. Association with serum gamma-glutamyltransferase levels and alcohol consumption on stroke and coronary artery disease: the Suita study. *Stroke* 42 (6): 1764-7, 2011.
- (140) Higashiyama A, et al. Alcohol consumption and cardiovascular disease incidence in men with and without hypertension: the Suita study. *Hypertens Res*; 36 (1): 58-64, 2013.
- (141) Naito M, Nakayama T, Okamura T, et al. Effect of a 4-year workplace-based physical activity intervention program on the blood lipid profiles of participating employees: The high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. *Atherosclerosis*; 197 (2): 784-790, 2008.
- (142) Subramanian S, Chait A. Hypertriglyceridemia secondary to obesity and diabetes. *Biochim Biophys Acta*. 1821: 819-25. 2012.
- (143) Aucott L, Gray D, Rothnie H, et al. Effects of lifestyle interventions and long-term weight loss on lipid outcomes - a systematic review. *Obes Rev*. 12: e412-25. 2011.
- (144) Eslick GD, Howe PR, Smith C, et al. Benefits of fish oil supplementation in hyperlipidemia: a systematic review and meta-analysis. *Int J Cardiol*. 136: 4-16. 2009.
- (145) Iso H, Kobayashi M, Ishihara J, et al. Intake of fish and n3 fatty acids and risk of coronary heart disease among Japanese: the Japan Public Health Center-Based (JPHC) Study Cohort I. *Circulation*. 113: 195-202. 2006.
- (146) Iso H, Sato S, Folsom AR, et al. Serum fatty acids and fish intake in rural Japanese, urban Japanese, Japanese American and Caucasian American men. *Int J Epidemiol*. 18: 374-81. 1989.
- (147) 日本動脈硬化学会編：動脈硬化性疾患予防ガイドライン2017年版. 日本動脈硬化学会, 2017.
- (148) 厚生労働省「日本人の食事摂取基準策定検討会」報告書(2015年度版).<http://www.mhlw.go.jp/stf/shingi/0000041824.html>
- (149) Dhingra R, Sullivan L, Jacques PF, et al. Soft drink consumption and risk of developing cardiometabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation*. 116: 480-8. 2007.
- (150) Hu FB, Willett WC. Optimal diets for prevention of coronary heart disease. *JAMA*. 288: 2569-78. 2002.

- (151) Ikehara S, Iso H, Toyoshima H, et al. Alcohol consumption and mortality from stroke and coronary heart disease among Japanese men and women: the Japan collaborative cohort study. *Stroke*. 39: 2936-42. 2008.
- (152) Lieber CS. Alcoholic fatty liver: its pathogenesis and mechanism of progression to inflammation and fibrosis. *Alcohol*. 34: 9-19. 2004.
- (153) Koba S, Tanaka H, Maruyama C, et al. Physical activity in the Japan population: association with blood lipid levels and effects in reducing cardiovascular and all-cause mortality. *J Atheroscler Thromb*. 18: 833-45. 2011.
- (154) Kelley GA, Kelley KS, Tran ZV. Walking, lipids, and lipoproteins: a meta-analysis of randomized controlled trials. *Prev Med*. 38: 651-61. 2004.
- (155) Kelley GA, Kelley KS. Impact of progressive resistance training on lipids and lipoproteins in adults: a meta-analysis of randomized controlled trials. *Prev Med*. 48: 9-19. 2009.
- (156) Sigal RJ, Kenny GP, Boule NG, et al. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med*. 147: 357-69. 2007.
- (157) Harada-Siba M, Arai H, Okumura T, Yokota K, Oikawa S, Nohara A, Okada T, Ohta T, Bujo H, Watanabe M, Wakatsuki A, Yamashita S. Multicenter study to determine the diagnosis criteria of heterozygous familial hypercholesterolemia in Japan. *J Atheroscler Thromb*. 2012; 19: 1019-26.
- (158) Lv J, Chen W, Sun D, et al. Gender-specific association between tobacco smoking and central obesity among 0.5 million Chinese people: the China Kadoorie Biobank Study. *PLoS One*. 2015 Apr 21; 10 (4): e0124586.
- (159) 永井, 坂根, 森谷ら. 朝食欠食, マクロニュートリエントバランスが若年健常者の食後血糖値, 満腹感, エネルギー消費量, および自律神経活動へ及ぼす影響. *糖尿病* 48 : 761-770, 2005.
- (160) Maruyama K, Sato S, Ohira T, et al. The joint impact on being overweight of self reported behaviours of eating quickly and eating until full: cross sectional survey. *BMJ*. 2008 Oct 21; 337: a2002.
- (161) Ma Y, Bertone ER, Stanek EJ, et al. Association between Eating Patterns and Obesity in a Free-living US Adult Population. *Am J Epidemiol* 2003; 158: 85-92.
- (162) Campion EW, Glynn RJ, DeLabry LO. Asymptomatic hyperuricemia. Risks and consequences in the Normative Aging Study. *Am J Med*. 1987; 82 (3): 421-6.
- (163) Richette P, Bardin T. Gout. *Lancet*. 2010; 375 (9711): 318-28.
- (164) Hall AP, Barry PE, Dawber TR, McNamara PM. Epidemiology of gout and hyperuricemia. A long-term population study. *Am J Med*. 1967; 42 (1): 27-37.
- (165) Yü T, Gutman AB. Uric acid nephrolithiasis in gout. Predisposing factors. *Ann Intern Med*. 1967; 67 (6): 1133-48.
- (166) Iseki K, Ikemiya Y, Inoue T, Iseki C, Kinjo K, Takishita S. Significance of hyperuricemia as a risk factor for developing ESRD in a screened cohort. *Am J Kidney Dis*. 2004; 44 (4): 642-50.
- (167) Weiner DE, Tighiouart H, Elsayed EF, Griffith JL, Salem DN, Levey AS. Uric acid and incident kidney disease in the community. *J Am Soc Nephrol*. 2008; 19 (6): 1204-11.
- (168) Choi HK, Liu S, Curhan G. Intake of purine-rich foods, protein, and dairy products and relationship to serum levels of uric acid: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum*. 2005; 52 (1): 283-9.

- (169) Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med.* 2004; 350 (11): 1093-103.
- (170) Choi JW, Ford ES, Gao X, Choi HK. Sugar-sweetened soft drinks, diet soft drinks, and serum uric acid level: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum.* 2008; 59 (1): 109-16.
- (171) Choi HK, Curhan G. Soft drinks, fructose consumption, and the risk of gout in men: prospective cohort study. *BMJ.* 2008; 336 (7639): 309-12.
- (172) Choi HK, Curhan G. Beer, liquor, and wine consumption and serum uric acid level: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum.* 2004; 51 (6): 1023-9.
- (173) Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Alcohol intake and risk of incident gout in men: a prospective study. *Lancet.* 2004; 363 (9417): 1277-81.
- (174) Matsuura F, Yamashita S, Nakamura T, Nishida M, Nozaki S, Funahashi T, Matsuzawa Y. Effect of visceral fat accumulation on uric acid metabolism in male obese subjects: visceral fat obesity is linked more closely to overproduction of uric acid than subcutaneous fat obesity. *Metabolism.* 1998; 47 (8): 929-33.
- (175) Choi HK, Atkinson K, Karlson EW, Curhan G. Obesity, weight change, hypertension, diuretic use, and risk of gout in men: the health professionals follow-up study. *Arch Intern Med.* 2005; 165 (7): 742-8.
- (176) Sadakane A, Tsutsumi A, Gotoh T, Ishikawa S, Ojima T, Kario K, Nakamura Y, Kayaba K. Dietary patterns and levels of blood pressure and serum lipids in a Japanese population. *J Epidemiol.* 2008; 18 (2): 58-67.
- (177) Petkevičienė J, Klumbienė J, Ramažauskienė V, Kriaucionienė V, Sakytė E, Grabauskas V. Diet and dyslipidemias in a Lithuanian rural population aged 25-64: the CINDI survey. *Medicina (Kaunas).* 2012; 48 (4): 211-7.
- (178) Dhingra R, Sullivan L, Jacques PF, Wang TJ, Fox CS, Meigs JB, D'Agostino RB, Gaziano JM, Vasan RS. Soft drink consumption and risk of developing cardiometabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation.* 2007; 116 (5): 480-8.
- (179) Wakabayashi I. Cross-sectional relationship between alcohol consumption and prevalence of metabolic syndrome in Japanese men and women. *J Atheroscler Thromb.* 2010; 17 (7): 695-704.
- (180) Oka R, Miura K, Sakurai M, Nakamura K, Yagi K, Miyamoto S, Moriuchi T, Mabuchi H, Koizumi J, Nomura H, Takeda Y, Inazu A, Nohara A, Kawashiri MA, Nagasawa S, Kobayashi J, Yamagishi M. Impacts of visceral adipose tissue and subcutaneous adipose tissue on metabolic risk factors in middle-aged Japanese. *Obesity (Silver Spring).* 2010; 18 (1): 153-60.
- (181) Choi HK, Ford ES, Li C, Curhan G. Prevalence of the metabolic syndrome in patients with gout: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum.* 2007; 57 (1): 109-15.
- (182) 斎田美穂. 高尿酸血症と生活習慣病, 糖代謝異常の関連に関する研究. 痛風と核酸代謝. 2000; 24 (2): 139-51.
- (183) 厚生労働省. 国民健康・栄養調査. http://www.mhlw.go.jp/bunya/kenkou/kenkou_eiou_chousa.html(アクセス2019年2月19日)
- (184) Zavaroni I, Mazza S, Fantuzzi M, Dall'Aglio E, Bonora E, Delsignore R, Passeri M,

- Reaven GM. Changes in insulin and lipid metabolism in males with asymptomatic hyperuricaemia. *J Intern Med.* 1993; 234 (1): 25-30.
- (185) Zavaroni I, Bonini L, Fantuzzi M, Dall'Aglio E, Passeri M, Reaven GM. Hyperinsulinaemia, obesity, and syndrome X. *J Intern Med.* 1994; 235 (1): 51-6.
- (186) Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, Fruchart JC, James WP, Loria CM, Smith SC Jr; International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; International Association for the Study of Obesity. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation.* 2009; 120 (16): 1640-5.
- (187) メタボリックシンドローム診断基準検討委員会. メタボリックシンドロームの定義と診断基準. 日本国内科学会雑誌. 2005; 94 (4): 794-809.
- (188) 日本痛風・核酸代謝学会ガイドライン改訂委員会. 高尿酸血症・痛風の治療ガイドライン(第2版). 大阪, メディカルレビュー社. 2010.
- (189) Benedict JD, Forsham PH, Stetten D Jr. The metabolism of uric acid in the normal and gouty human studied with the aid of isotopic uric acid. *J Biol Chem.* 1949; 181 (1): 183-93.
- (190) Scott JT, Holloway VP, Glass HI, Arnot RN. Studies of uric acid pool size and turnover rate. *Ann Rheum Dis.* 1969; 28 (4): 366-73.
- (191) Ryu S, Chang Y, Zhang Y, Kim SG, Cho J, Son HJ, Shin H, Guallar E. A cohort study of hyperuricemia in middle-aged South Korean men. *Am J Epidemiol.* 2012; 175 (2): 133-43.
- (192) Nakamura K, Sakurai M, Miura K, Morikawa Y, Nagasawa SY, Ishizaki M, Kido T, Naruse Y, Nakashima M, Nogawa K, Suwazono Y, Nakagawa H. HOMA-IR and the risk of hyperuricemia: a prospective study in non-diabetic Japanese men. *Diabetes Res Clin Pract.* 2014; 106 (1): 154-60.
- (193) Leyva F, Wingrove CS, Godsland IF, Stevenson JC. The glycolytic pathway to coronary heart disease: a hypothesis. *Metabolism.* 1998; 47 (6): 657-62.
- (194) Facchini F, Chen YD, Hollenbeck CB, Reaven GM. Relationship between resistance to insulin-mediated glucose uptake, urinary uric acid clearance, and plasma uric acid concentration. *JAMA.* 1991; 266 (21): 3008-11.
- (195) Quiñones Galvan A, Natali A, Baldi S, Frascerra S, Sanna G, Ciociaro D, Ferrannini E. Effect of insulin on uric acid excretion in humans. *Am J Physiol.* 1995; 268 (1 Pt 1): E1-5.
- (196) Hakoda M, Masunari N, Yamada M, Fujiwara S, Suzuki G, Kodama K, Kasagi F. Serum uric acid concentration as a risk factor for cardiovascular mortality: a longterm cohort study of atomic bomb survivors. *J Rheumatol.* 2005; 32 (5): 906-12.
- (197) Fang J, Alderman MH. Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971-1992. National Health and Nutrition Examination Survey. *JAMA.* 2000; 283 (18): 2404-10.
- (198) Puddu PE, Lanti M, Menotti A, Mancini M, Zanchetti A, Cirillo M, Angeletti M, Panarelli W; Gubbio Study Research Group. Serum uric acid for short-term prediction of cardiovascular disease incidence in the Gubbio population Study. *Acta Cardiol.* 2001; 56 (4): 243-51.

- (199) Bos MJ, Koudstaal PJ, Hofman A, Witteman JC, Breteler MM. Uric acid is a risk factor for myocardial infarction and stroke: the Rotterdam study. *Stroke*. 2006;37(6):1503-7.
- (200) Strasak A, Ruttmann E, Brant L, Kelleher C, Klenk J, Concin H, Diem G, Pfeiffer K, Ulmer H; VHM&PP Study Group. Serum uric acid and risk of cardiovascular mortality: a prospective long-term study of 83,683 Austrian men. *Clin Chem*. 2008; 54 (2): 273-84.
- (201) Strasak AM, Kelleher CC, Brant LJ, Rapp K, Ruttmann E, Concin H, Diem G, Pfeiffer KP, Ulmer H; VHM&PP Study Group. Serum uric acid is an independent predictor for all major forms of cardiovascular death in 28,613 elderly women: a prospective 21-year follow-up study. *Int J Cardiol*. 2008; 125 (2): 232-9.
- (202) Chien KL, Hsu HC, Sung FC, Su TC, Chen MF, Lee YT. Hyperuricemia as a risk factor on cardiovascular events in Taiwan: The Chin-Shan Community Cardiovascular Cohort Study. *Atherosclerosis*. 2005; 183 (1): 147-55.
- (203) Culleton BF, Larson MG, Kannel WB, Levy D. Serum uric acid and risk for cardiovascular disease and death: the Framingham Heart Study. *Ann Intern Med*. 1999; 131 (1): 7-13.
- (204) Wannamethee SG, Shaper AG, Whincup PH. Serum urate and the risk of major coronary heart disease events. *Heart*. 1997; 78 (2): 147-53.
- (205) Jee SH, Lee SY, Kim MT. Serum uric acid and risk of death from cancer, cardiovascular disease or all causes in men. *Eur J Cardiovasc Prev Rehabil*. 2004; 11 (3): 185-91.
- (206) Sakata K, Hashimoto T, Ueshima H, Okayama A; NIPPON DATA 80 Research Group. Absence of an association between serum uric acid and mortality from cardiovascular disease: NIPPON DATA 80, 1980-1994. National Integrated Projects for Prospective Observation of Non-communicable Diseases and its Trend in the Aged. *Eur J Epidemiol*. 2001; 17 (5): 461-8.
- (207) 箱田雅之. 血清尿酸値と心血管リスク. 痛風と核酸代謝. 2008; 32 (1): 1-12.
- (208) Kim SY, Guevara JP, Kim KM, Choi HK, Heitjan DF, Albert DA. Hyperuricemia and coronary heart disease: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken)*. 2010; 62 (2): 170-80.
- (209) Kim SY, Guevara JP, Kim KM, Choi HK, Heitjan DF, Albert DA. Hyperuricemia and risk of stroke: a systematic review and meta-analysis. *Arthritis Rheum*. 2009; 61 (7): 885-92.
- (210) 富田眞佐子, 横田和彦, 水野正一. 健康診断, 人間ドックにおける尿酸測定の意義. 高尿酸血症と痛風. 2010; 18 (1): 67-71.
- (211) Yu TS, Berger L, Gutman AB. Renal function in gout. II. Effect of uric acid loading on renal excretion of uric acid. *Am J Med*. 1962; 33: 829-44.
- (212) Yamamoto T, Moriwaki Y, Takahashi S, Yamakita J, Tsutsumi Z, Ohata H, Hiroishi K, Nakano T, Higashino K. Effect of ethanol and fructose on plasma uridine and purine bases. *Metabolism*. 1997; 46 (5): 544-7.
- (213) Yamamoto T, Moriwaki Y, Takahashi S, Tsutsumi Z, Yamakita J, Higashino K. Effects of fructose and xylitol on the urinary excretion of adenosine, uridine, and purine bases. *Metabolism*. 1999; 48 (4): 520-4.
- (214) Villegas R, Xiang YB, Elasy T, Xu WH, Cai H, Cai Q, Linton MF, Fazio S, Zheng W, Shu XO. Purine-rich foods, protein intake, and the prevalence of hyperuricemia: the Shanghai Men's Health Study. *Nutr Metab Cardiovasc Dis*. 2012; 22 (5): 409-16.
- (215) Gao X, Qi L, Qiao N, Choi HK, Curhan G, Tucker KL, Ascherio A. Intake of added sugar

- and sugar-sweetened drink and serum uric acid concentration in US men and women. *Hypertension*. 2007; 50 (2): 306-12.
- (216) Dhingra R, Sullivan L, Jacques PF, Wang TJ, Fox CS, Meigs JB, D'Agostino RB, Gaziano JM, Vasan RS. Soft drink consumption and risk of developing cardiometabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation*. 2007; 116 (5): 480-8.
- (217) Bes-Rastrollo M, Sánchez-Villegas A, Gómez-Gracia E, Martínez JA, Pajares RM, Martínez-González MA. Predictors of weight gain in a Mediterranean cohort: the Seguimiento Universidad de Navarra Study 1. *Am J Clin Nutr*. 2006; 83 (2): 362-70.
- (218) Yamamoto T, Moriwaki Y, Takahashi S, Tsutsumi Z, Yamakita J, Nakano T, Higashino K. Effect of glucose on the plasma concentration and urinary excretion of uridine and purine bases. *Metabolism*. 1999; 48 (3): 338-41.
- (219) Kilinenberg JR, Goldfinger SE, Seegmiller JE. The effectiveness of the xanthine oxidase inhibitor allopurinol in the treatment of gout. *Ann Intern Med*. 1965; 62: 639-47.
- (220) Siener R, Hesse A. The effect of a vegetarian and different omnivorous diets on urinary risk factors for uric acid stone formation. *Eur J Nutr*. 2003; 42 (6): 332-7.
- (221) Zhou BF, Stamler J, Dennis B, Moag-Stahlberg A, Okuda N, Robertson C, Zhao L, Chan Q, Elliott P; INTERMAP Research Group. Nutrient intakes of middle-aged men and women in China, Japan, United Kingdom, and United States in the late 1990s: the INTERMAP study. *J Hum Hypertens*. 2003; 17 (9): 623-30.
- (222) 国税庁. 酒税 お酒に関する情報 酒税課税関係統計資料. <http://www.nta.go.jp/shiraberu/senmonjoho/sake/tokei/mokuji.htm>(アクセス2013年2月22日)
- (223) Kaneko K, Yamanobe T, Fujimori S. Determination of purine contents of alcoholic beverages using high performance liquid chromatography. *Biomed Chromatogr*. 2009; 23 (8): 858-64.
- (224) Yamamoto T, Moriwaki Y, Takahashi S, Tsutsumi Z, Ka T, Fukuchi M, Hada T. Effect of beer on the plasma concentrations of uridine and purine bases. *Metabolism*. 2002; 51 (10): 1317-23.
- (225) Yamamoto T, Moriwaki Y, Ka T, Inokuchi T, Takahashi S, Tsutsumi Z, Fukuchi M, Hada T. Effect of purine-free low-malt liquor (happo-shu) on the plasma concentrations and urinary excretion of purine bases and uridine--comparison between purine-free and regular happo-shu. *Horm Metab Res*. 2004; 36 (4): 231-7.
- (226) Nishimura T, Shimizu T, Mineo I, Kawachi M, Ono A, Nakajima H, Kuwajima M, Kono N, Matsuzawa Y. Influence of daily drinking habits on ethanol-induced hyperuricemia. *Metabolism*. 1994; 43 (6): 745-8.
- (227) 清水孝郎, 西村隆通, 大野昭. 日常飲酒量が飲酒後の高尿酸血症に及ぼす影響. プリン・ピリミジン代謝. 1992; 16 (1): 9-13.
- (228) Yamanaka H, Kamatani N, Hakoda M, Terai C, Kawaguchi R, Kashiwazaki S. Analysis of the genotypes for aldehyde dehydrogenase 2 in Japanese patients with primary gout. *Adv Exp Med Biol*. 1994; 370: 53-6.
- (229) Faller J, Fox IH. Ethanol-induced hyperuricemia: evidence for increased urate production by activation of adenine nucleotide turnover. *N Engl J Med*. 1982; 307 (26): 1598-602.
- (230) Nakamura K, Sakurai M, Miura K, Morikawa Y, Yoshita K, Ishizaki M, Kido T, Naruse Y, Suwazono Y, Nakagawa H. Alcohol intake and the risk of hyperuricaemia: a 6-year

- prospective study in Japanese men. *Nutr Metab Cardiovasc Dis.* 2012; 22 (11): 989-96.
- (231) Yamashita S, Matsuzawa Y, Tokunaga K, Fujioka S, Tarui S. Studies on the impaired metabolism of uric acid in obese subjects: marked reduction of renal urate excretion and its improvement by a low-calorie diet. *Int J Obes.* 1986; 10 (4): 255-64.
- (232) Stewart KJ, Bacher AC, Turner K, Lim JG, Hees PS, Shapiro EP, Tayback M, Ouyang P. Exercise and risk factors associated with metabolic syndrome in older adults. *Am J Prev Med.* 2005 Jan; 28 (1): 9-18.
- (233) Katzel LI, Bleecker ER, Rogus EM, Goldberg AP. Sequential effects of aerobic exercise training and weight loss on risk factors for coronary disease in healthy, obese middle-aged and older men. *Metabolism.* 1997; 46 (12): 1441-7.
- (234) Yamanaka H, Kawagoe Y, Taniguchi A, Kaneko N, Kimata S, Hosoda S, Kamatani N, Kashiwazaki S. Accelerated purine nucleotide degradation by anaerobic but not by aerobic ergometer muscle exercise. *Metabolism.* 1992; 41 (4): 364-9.
- (235) Kaya M, Moriwaki Y, Ka T, Inokuchi T, Yamamoto A, Takahashi S, Tsutsumi Z, Tsuzita J, Oku Y, Yamamoto T. Plasma concentrations and urinary excretion of purine bases (uric acid, hypoxanthine, and xanthine) and oxypurinol after rigorous exercise. *Metabolism.* 2006; 55 (1): 103-7.
- (236) 伊藤朗, 三上俊夫, 丹信介. 各種運動時の血清尿酸値の動態. 尿酸. 1984; 8 (1): 38-47.
- (237) Ka T, Yamamoto T, Moriwaki Y, Kaya M, Tsujita J, Takahashi S, Tsutsumi Z, Fukuchi M, Hada T. Effect of exercise and beer on the plasma concentration and urinary excretion of purine bases. *J Rheumatol.* 2003; 30 (5): 1036-42.
- (238) Ohno M, Ka T, Inokuchi T, Moriwaki Y, Yamamoto A, Takahashi S, Tsutsumi Z, Tsuzita J, Yamamoto T, Nishiguchi S. Effects of exercise and grape juice ingestion in combination on plasma concentrations of purine bases and uridine. *Clin Chim Acta.* 2008; 388 (1-2): 167-72.
- (239) Seegmiller JE, Rosenbloom FM, Kelley WN. Enzyme defect associated with a sex-linked human neurological disorder and excessive purine synthesis. *Science.* 1967; 155 (3770): 1682-4.
- (240) Kelley WN, Rosenbloom FM, Henderson JF, Seegmiller JE. A specific enzyme defect in gout associated with overproduction of uric acid. *Proc Natl Acad Sci U S A.* 1967; 57 (6): 1735-9.
- (241) Puig JG, Miranda ME, Mateos FA, Picazo ML, Jiménez ML, Calvin TS, Gil AA. Hereditary nephropathy associated with hyperuricemia and gout. *Arch Intern Med.* 1993; 153 (3): 357-65.
- (242) Pencina MJ, et al. *Circulation* 2009; 119: 3078-3084.
- (243) NIPPON DATA80 Research Group. *Circ J* 2006; 70: 1249-1255.
- (244) 上島弘嗣編著,「NIPPON DATAからみた循環器疾患のエビデンス」,日本医事新報社.
- (245) Nishimura K, Okamura T, Watanabe M, et al. Predicting coronary heart disease using risk factor categories for a Japanese urban population, and comparison with the framingham risk score: the suita study. *J Atheroscler Thromb.* 2014;21(8):784-98.
- (246) The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice. *European Heart Journal* 2012; 33: 1635-1701.
- (247) Arima H, et al. *Hypertens Res* 2009; 32: 1119-1122.
- (248) Ishikawa S, et al.; Jichi Medical School (JMS) Cohort Study Group. *J Epidemiol.* 2009; 19:

- 101-106.
- (249) Matsumoto M, et al.; Jichi Medical School (JMS) Cohort Study Group. *J Epidemiol*. 2009; 19: 94-100.
- (250) Tanabe N, et al.; The Japan Arteriosclerosis Longitudinal Study Group. *Circ J*. 2010; 74: 1346-1356.
- (251) Kokubo Y, et al. *Hypertens Res* 2008; 31: 2027-2035.
- (252) Alberti KGMM, et al. *Circulation* 2009; 120: 1640-1645.
- (253) メタボリックシンドローム診断基準検討委員会 日本内科学会雑誌2005 ; 94: 749.
- (254) Barker DJ, et al. *Diabetologia* 1993; 36: 62-67.
- (255) Curhan GC, et al. *Circulation* 1996; 94: 3246-3250.
- (256) Miura K, et al. *Am J Epidemiol* 2001; 153: 783-789.
- (257) Nanri A, et al. *J Epidemiol Community Health* 2011; 65: 1104-1110.
- (258) Chou WT, et al. *Circ J* 2012, 2013; 77: 679-686.
- (259) Go, A.S., et al., Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med*, 2004. 351(13): p. 1296-305.
- (260) Levey, A.S., et al., The definition, classification, and prognosis of chronic kidney disease: a KDIGO Controversies Conference report. *Kidney Int*, 2011. 80(1): p. 17-28.
- (261) Nagai, K., et al., Relationship between Estimated Glomerular Filtration Rate and Cardiovascular Mortality in a Japanese Cohort with Long-Term Follow-Up. *PLoS One*, 2016. 11(6): p. e0156792.
- (262) わが国の慢性透析療法の現況(2016年12月31日現在). 2018, 日本透析医学会統計調査委員会.
- (263) Pugliese, G., Updating the natural history of diabetic nephropathy. *Acta Diabetol*, 2014. 51(6): p. 905-15.
- (264) Bhan, I. and R. Thadhani, Calsium, phosphorus, and vitamin D in kidney disease, in *Handbook of nutrition and the kidney*, W.E. Mitch and T.A. Ikizler, Editors. 2010, Wolter Kluwer: Philadelphia. p. 50-57.
- (265) 厚生労働省. 人口動態調査 2017 [Available from: <https://www.e-stat.go.jp/stat-search/files?page=1&query=%E4%BA%BA%E5%8F%A3%E5%8B%95%E6%85%8B%E8%AA%BF%E6%9F%BB&layout=dataset&lid=000001101884>] accessed March 13 2018.
- (266) Sasaki N. The Relationship of Salt Intake to Hypertension in the Japanese. *Geriatrics* 1964; 19: 735-44. [published Online First: 1964/10/01]
- (267) 厚生労働省. 平成28年国民健康・栄養調査報告 2017 [Available from: <http://www.mhlw.go.jp/bunya/kenkou/eiyu/h28-houkoku.html>] accessed March 9 2018.
- (268) 奥田奈賀子・岡山明. ナトリウム・カリウム摂取と循環器疾患. *循環器内科* 2011; 70 (6): 546-54.
- (269) He FJ, Li J, Macgregor GA. Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ* 2013;346:f1325. doi: 10.1136/bmj.f1325 [published Online First: 2013/04/06]
- (270) Cook NR, Cutler JA, Obarzanek E, et al. Long term effects of dietary sodium reduction on cardiovascular disease outcomes: observational follow-up of the trials of hypertension prevention (TOHP). *BMJ* 2007; 334 (7599): 885-8. doi: 10.1136/bmj.39147.604896.55 [published Online First: 2007/04/24]
- (271) Aburto NJ, Hanson S, Gutierrez H, et al. Effect of increased potassium intake on cardiovascular risk factors and disease: systematic review and meta-analyses. *BMJ* 2013; 346: f1378. doi: 10.1136/bmj.f1378 [published Online First: 2013/04/06]
- (272) Jackson SL, Cogswell ME, Zhao L, et al. Association Between Urinary Sodium and

- Potassium Excretion and Blood Pressure Among Adults in the United States: National Health and Nutrition Examination Survey, 2014. *Circulation* 2018; 137 (3): 237-46. doi: 10.1161/CIRCULATIONAHA.117.029193.
- (273) Okayama A, Okuda N, Miura K, et al. Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA 80 cohort study. *BMJ open* 2016; 6 (7): e011632. doi: 10.1136/bmjopen-2016-011632.
 - (274) Chang HY, Hu YW, Yue CS, et al. Effect of potassium-enriched salt on cardiovascular mortality and medical expenses of elderly men. *The American journal of clinical nutrition* 2006; 83 (6): 1289-96. [published Online First: 2006/06/10]
 - (275) Harvard Report on Cancer Prevention. Volume 1: Causes of human cancer. *Cancer causes & control: CCC*. 1996;7 Suppl 1:S3-59.
 - (276) Tsugane S. Salt, salted food intake, and risk of gastric cancer: epidemiologic evidence. *Cancer science*. 2005; 96 (1):1-6.
 - (277) 塚本徹哉, 立松正衛. 基礎医学から 高食塩食とピロリ菌による発癌. *日本医事新報*. 2005 (4252): 11-13.
 - (278) 国立がん研究センターがん対策情報センター. がん登録・統計グラフデータベース. 2018; http://gdb.ganjoho.jp/graph_db/index. Accessed March 13, 2018.
 - (279) Shikata K, Kiyohara Y, Kubo M, et al. A prospective study of dietary salt intake and gastric cancer incidence in a defined Japanese population: the Hisayama study. *International journal of cancer Journal international du cancer*. 2006; 119 (1): 196-201.
 - (280) 黒川清. 食塩と高血圧. *日病態栄養会誌* 2001; 5 (1-6): 18.
 - (281) 奥田奈賀子 上島弘嗣. 海外における食塩摂取量の推移と現状—24時間尿を用いた推定食塩摂取量より—. *カレントテラピー* 2013; 31 (10): 14-19.
 - (282) He FJ, Brinsden HC, MacGregor GA. Salt reduction in the United Kingdom: a successful experiment in public health. *Journal of human hypertension* 2014; 28 (6): 345-52. doi: 10.1038/jhh.2013.105.
 - (283) Bernstein AM, Willett WC. Trends in 24-h urinary sodium excretion in the United States, 1957-2003: a systematic review. *The American journal of clinical nutrition* 2010; 92 (5): 1172-80. doi: 10.3945/ajcn.2010.29367 [published Online First: 2010/09/10]
 - (284) Sasaki N. The Relationship of Salt Intake to Hypertension in the Japanese. *Geriatrics* 1964; 19: 735-44. [published Online First: 1964/10/01]
 - (285) Uechi K, Sugimoto M, Kobayashi S, et al. Urine 24-Hour Sodium Excretion Decreased between 1953 and 2014 in Japan, but Estimated Intake Still Exceeds the WHO Recommendation. *The Journal of nutrition* 2017; 147 (3): 390-97. doi: 10.3945/jn.116.240960.
 - (286) Group ICR. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. Intersalt Cooperative Research Group. *BMJ* 1988; 297 (6644): 319-28. [published Online First: 1988/07/30]
 - (287) Zhou BF, Stamler J, Dennis B, et al. Nutrient intakes of middle-aged men and women in China, Japan, United Kingdom, and United States in the late 1990s: the INTERMAP study. *Journal of human hypertension* 2003; 17 (9): 623-30. doi: 10.1038/sj.jhh.1001605 [published Online First: 2003/09/19]
 - (288) Dauchet L, Amouyel P, Hercberg S, Dallongeville J. Fruit and vegetable consumption and risk of coronary heart disease: a meta-analysis of cohort studies. *The Journal of nutrition*. 2006; 136 (10): 2588-2593.
 - (289) Okuda N, Miura K, Okayama A, et al. Fruit and vegetable intake and mortality from cardiovascular disease in Japan: a 24-year follow-up of the NIPPON DATA80 Study.

- European journal of clinical nutrition. 2015; 69 (4): 482-488.
- (290) 厚生労働省. 平成28年国民健康・栄養調査報告. 2017; <http://www.mhlw.go.jp/bunya/kenkou/eiyou/h28-houkoku.html>. Accessed March 9, 2018.
- (291) Institute NC. Usual Dietary Intakes: Food Intakes, US Population, 2001-04. Risk Factor Monitoring and Methods Branch Web site. Applied Research Program. <https://epi.grants.cancer.gov/diet/usualintakes/pop/2001-04/>. Accessed March 1, 2018.
- (292) 厚生省公衆衛生局栄養課. 国民栄養の現状(昭和22年国民栄養調査成績) 1948 [Available from: http://www.nibiohn.go.jp/eiken/chosa/kokumin_eiyou/doc_year/1947/1947_kek.pdf] accessed March 1st 2018.
- (293) Okuda N, Okayama A, Miura K, et al. Food sources of dietary sodium in the Japanese adult population: the international study of macro-/micronutrients and blood pressure (INTERMAP). European journal of nutrition 2017; 56 (3): 1269-80. doi: 10.1007/s00394-016-1177-1.
- (294) 堀田宗徳. 最近の中食の動向. 日本調理科学会誌 2007; 40 (2): 104-08.
- (295) 村本信幸. 外食産業市場規模と外食比率. 日本調理科学会誌 1995; 28 (4): 290-96.
- (296) 独立行政法人農畜産業振興機構. 平成28年度カット野菜・冷凍野菜・野菜惣菜に係る小売販売動向調査. 2017.
- (297) Ikeda N, Okuda N, Tsubota-Utsugi M, et al. Association of Energy Intake With the Lack of in-Person Review of Household Dietary Records: Analysis of Japan National Health and Nutrition Surveys From 1997 to 2011. Journal of epidemiology / Japan Epidemiological Association 2016; 26 (2): 84-91. doi: 10.2188/jea.JE20150048.
- (298) 厚生労働省. 介護予防マニュアル(改訂版) 2012 [Available from: <http://www.mhlw.go.jp/topics/2009/05/tp0501-1.html>] accessed March 2 2018.
- (299) 葛谷雅文. 総論. 静脈経腸栄養. 2007; 22 (4): 433-7.
- (300) Eng PM, Kawachi I, Fitzmaurice G, et al. Effects of marital transitions on changes in dietary and other health behaviours in US male health professionals. Journal of epidemiology and community health 2005; 59 (1): 56-62. doi: 10.1136/jech.2004.020073.
- (301) Nowson C, O'Connell S. Protein Requirements and Recommendations for Older People: A Review. Nutrients 2015; 7 (8): 6874-99. doi: 10.3390/nu7085311.
- (302) Kobayashi S, Asakura K, Suga H, et al. High protein intake is associated with low prevalence of frailty among old Japanese women: a multicenter cross-sectional study. Nutrition journal 2013; 12: 164. doi: 10.1186/1475-2891-12-164.
- (303) 厚生労働省. 日本人の食事摂取基準2015年 2014 [Available from: <http://www.mhlw.go.jp/stf/shingi/0000041824.html>] accessed March 12 2018.
- (304) 厚生労働省健康局健康課<<http://www.mhlw.go.jp/>> : 平成28年国民健康・栄養調査結果
- (305) Physical Activity Guidelines Advisory Committee: Physical Activity Guidelines Advisory Committee Report. 2008. U. S. Department of Health and Human Services, 2008.
- (306) 日本動脈硬化学会編:動脈硬化性疾患予防ガイドライン2017年版. 日本動脈硬化学会, 2017.
- (307) 日本高血圧学会高血圧治療ガイドライン作成委員会編. 高血圧治療ガイドライン2019. 日本高血圧学会, 2019.
- (308) Lee IM, et al.: Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. Lancet, 380: 219-229, 2012.
- (309) Sasaki T, et al.: Role of AMPK and PPAR γ 1 in exercise-induced lipoprotein lipase in skeletal muscle. Am J Physiol Endocrinol Metab, 306: E1085-E1092, 2014.
- (310) Kodama S, et al.: Effect of aerobic exercise training on serum levels of high-density

- lipoprotein cholesterol: a meta-analysis. Arch Intern Med, 167: 999-1008, 2007.
- (311) Kraus WE, et al.: Effects of the amount and intensity of exercise on plasma lipoproteins. N Engl J Med, 347: 1483-1492, 2002.
 - (312) WHO: Recommended levels of physical activity for adults aged 18-64 years. <http://www.who.int/dietphysicalactivity/factsheet_adults/en/>
 - (313) Weiss CO: Frailty and chronic diseases in older adults. Clin Geriatr Med, 27: 39-52, 2011.
 - (314) Walston J, et al.: Research agenda for frailty in older adults: toward a better understanding of physiology and etiology: summary from the American Geriatrics Society National Institute on Aging Research Conference on Frailty in Older Adults. J Am Geriatr Soc, 54: 991-1001, 2006.
 - (315) Binder EF, et al.: Effects of exercise training on frailty in community-dwelling older adults: results of a randomized, controlled trial. J Am Geriatr Soc, 50: 1921-1928, 2002.
 - (316) Gitlin LN, et al.: A randomized trial of a multicomponent home intervention to reduce functional difficulties in older adults. J Am Geriatr Soc, 54: 809-816, 2006.
 - (317) 介護予防マニュアル改訂委員会:「第3章運動器の機能向上マニュアル」介護予防マニュアル改訂版, p46-64, 平成24年3月, 三菱総合研究所人間・生活研究本部, 東京. <http://www.mhlw.go.jp/topics/2009/05/dl/tp0501-1_1.pdf>
 - (318) Peterson MJ, et al.: Physical activity as a preventative factor for frailty: the health, aging, and body composition study. J Gerontol A Biol Sci Med Sci, 64: 61-68, 2009.
 - (319) WHO: Global Recommendations on Physical Activity for Health. 2010. <http://whqlibdoc.who.int/publications/2010/9789241599979_eng.pdf>
 - (320) Ikeda N, et al.: Adult mortality attributable to preventable risk factors for non-communicable diseases and injuries in Japan: a comparative risk assessment. PLoS Med, 9: e1001160, 2012.
 - (321) Sedentary Behavior Research Network: Standardized use of the terms "sedentary" and "sedentary behaviors". Appl Physiol Nutr Metab, 37: 540-542, 2012.
 - (322) Bauman A, et al.: The descriptive epidemiology of sitting. A 20-country comparison using the International Physical Activity Questionnaire (IPAQ). Am J Prev Med, 41: 228-235, 2011.
 - (323) Thorp A, et al.: Sedentary Behaviors and subsequent health outcomes: a systematic review of longitudinal studies, 1996-2011. Am J Prev Med, 41: 207-215, 2011.
 - (324) van der Ploeg HP, et al.: Sitting time and all-cause mortality risk in 222,497 Australian adults. Arch Intern Med, 172: 494-500, 2012.
 - (325) Honda T, et al.: Sedentary bout durations and metabolic syndrome among working adults: a prospective cohort study. BMC Public Health, 16: 888, 2016.
 - (326) Dunstan DW, et al.: Too much sitting-a health hazard. Diabetes Res Clin Pract, 97: 368-376, 2012.
 - (327) Armstrong MEG, et al.: Frequent physical activity may not reduce vascular disease risk as much as moderate activity. Circulation, 131: 721-729, 2015.
 - (328) Hirayama T. Non-smoking wives of heavy smokers have a higher risk of lung cancer: a study from Japan. BMJ 1981; 282:183-185.
 - (329) Fischer F, Kraemer A. Meta-analysis of the association between second-hand smoke exposure and ischaemic heart diseases, COPD and stroke. BMC Public Health 2015; 15:1202 DOI 10.1186/s12889-015-2489-4.
 - (330) 外務省. 水銀に関する水俣条約. https://www.mofa.go.jp/mofaj/ila/et/page22_001867.html
 - (331) 日本高血圧学会高血圧治療ガイドライン作成委員会編. 高血圧治療ガイドライン2019. 日本高血圧学会,

2019.

- (332) Mancia G, et al. 2007 Guidelines for the Management of Arterial Hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens.* 2007; 25: 1105-1187.
- (333) Chobanian AV, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA.* 2003; 289: 2560-2572.
- (334) Pickering TG, et al. Recommendations for blood pressure measurement in humans and experimental animals: Part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension.* 2005; 45: 142-161.
- (335) O'Brien E, et al. European Society of Hypertension recommendations for conventional, ambulatory and home blood pressure measurement. *J Hypertens.* 2003; 21: 821-848.
- (336) 日本循環器学会. 24時間血圧計の使用(ABPM)基準に関するガイドライン(2010年改訂版). http://www.j-circ.or.jp/guideline/pdf/JCS2010_shimada_h.pdf
- (337) Hozawa A, et al. Prognosis of isolated systolic and isolated diastolic hypertension as assessed by self-measurement of blood pressure at home: the Ohasama study. *Arch Intern Med.* 2000; 160: 3301-3306.
- (338) Imai Y, et al. Usefulness of home blood pressure measurements in assessing the effect of treatment in a single-blind placebo-controlled open trial. *J Hypertens.* 2001; 19: 179-185.
- (339) Iwahori T, Miura K, Obayashi K et al. Seasonal variation in home blood pressure: findings from nationwide web-based monitoring in Japan. *BMJ Open* 2018; 8: e017351.
- (340) Mancia G, et al. Ambulatory blood pressure is superior to clinic blood pressure in predicting treatment-induced regression of left ventricular hypertrophy. SAMPLE Study Group. *Study on Ambulatory Monitoring of Blood Pressure and Lisinopril Evaluation. Circulation.* 1997; 95: 1464-1470.
- (341) Staessen JA, et al. Predicting cardiovascular risk using conventional vs ambulatory blood pressure in older patients with systolic hypertension. *Systolic Hypertension in Europe Trial Investigators. JAMA.* 1999; 282: 539-546.
- (342) O'Brien E, Parati G, Stergiou G et al. European Society of Hypertension position paper on ambulatory blood pressure monitoring. *J Hypertens* 2013; 31: 1731-68.
- (343) Whelton PK, Carey RM, Aronow WS et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension* 2018; 71: e13-e115.
- (344) Asayama K, Ohkubo T, Yoshida S et al. Stroke risk and antihypertensive drug treatment in the general population: the Japan arteriosclerosis longitudinal study. *J Hypertens* 2009; 27: 357-64.
- (345) Staessen JA, Thijs L, Ohkubo T et al. Thirty years of research on diagnostic and therapeutic thresholds for the self-measured blood pressure at home. *Blood Press Monit* 2008; 13: 352-65.

- (346) Niranen TJ, Asayama K, Thijs L et al. Outcome-driven thresholds for home blood pressure measurement: international database of home blood pressure in relation to cardiovascular outcome. *Hypertension* 2013; 61: 27-34.
- (347) Ettehad D, Emdin CA, Kiran A et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 2016; 387: 957-967.
- (348) INTERSALT Cooperative Research Group. INTERSALT: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. *Br Med J* 1998; 297: 319-328.
- (349) Kuft FC, Fineberg NS, Sloan RS. Estimating dietary sodium intake in individuals receiving a randomly fluctuating intake. *Hypertension* 1982; 4: 805-808.
- (350) Yamasue K, Tochikubo O, Kono E, et al. Self-monitoring of home blood pressure with estimation of daily salt intake using a new electrical device. *J Hum Hypertens* 2006; 20: 593-598.
- (351) 川崎晃一, 上園慶子, 伊藤和枝, 上野道雄, 藤島正敏. 尿中クレアチニン排泄量予測値と起床後2回目のスポット尿を用いた24時間尿中ナトリウムならびにカリウム排泄量の推定法. *健康科学* 1988; 10: 115-120.
- (352) Tanaka T, Okamura T, Miura K, et al. A simple method to estimate populational 24-h urinary sodium and potassium excretion using a casual urine specimen. *J Hum Hypertens* 2002; 16: 97-103.
- (353) Yoshita K, Miura K, Okayama A, et al. A validation study on food composition tables for the international cooperative INTERMAP study in Japan. *Environ Health Prev Med* 2005; 10: 150-156.
- (354) Anderson CA, Appel LJ, Okuda N et al. Dietary sources of sodium in China, Japan, the United Kingdom, and the United States, women and men aged 40 to 59 years: the INTERMAP study. *J Am Diet Assoc*. 2010; 110: 736-745.
- (355) Sasaki S, Yanagimori R, Amano K. Validity of a self-administered diet history questionnaire for assessment of sodium and potassium - Comparison with single 24-hour urinary excretion. *Jpn Circ J* 1998; 62: 431-435.
- (356) Kobayashi S, Honda S, Murakami K, et al. Both comprehensive and brief self-administered diet history questionnaires satisfactorily rank nutrient intakes in Japanese adults. *J Epidemiol* 2012; 22: 151-159.
- (357) 日本高血圧学会減塩委員会. 日本高血圧学会減塩委員会報告2012. 日本高血圧学会; 2012.
- (358) 糖尿病診断基準に関する調査検討委員会：糖尿病の分類と診断基準に関する委員会報告(国際標準化対応版)糖尿病55(7) : 485-504, 2012.
- (359) World Health Organization: Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia. Report of a WHO/IDF consultation, 2006 (http://www.who.int/diabetes/publications/diagnosis_diabetes2006/en/)
- (360) 日本糖尿病学会：熊本宣言2013(<http://www.jds.or.jp/>)
- (361) 動脈硬化性疾患予防ガイドライン2012年版. 日本動脈硬化学会, 2012.
- (362) Nakamura M, Sato S, Shimamoto T. Improvement in Japanese clinical laboratory measurements of total cholesterol and HDL-cholesterol by the US Cholesterol Reference Method Laboratory Network. *J Atheroscler Thromb*. 10: 145-53. 2003.
- (363) Miida T, Nishimura K, Okamura T, et al. A multicenter study on the precision and accuracy of homogeneous assays for LDL-cholesterol: comparison with a beta-quantification method using fresh serum obtained from non-diseased and diseased

subjects. *Atherosclerosis*. 225: 208-15. 2012.

- (364) Miida T, et al. Homogeneous Assays for LDL-C and HDL-C are Reliable in Both the Postprandial and Fasting State. *J Atheroscler Thromb.* 2017; 24 (6): 583-599.
- (365) Sugimoto K, Isobe K, Kawakami Y, Yamada N. The relationship between non-HDL cholesterol and other lipid parameters in Japanese subjects. *J Atheroscler Thromb.* 12: 107-10. 2005.
- (366) Shimano H, Arai H, Harada-Shiba M, et al. Proposed guidelines for hypertriglyceridemia in Japan with non-HDL cholesterol as the second target. *J Atheroscler Thromb.* 15: 116-21. 2008.
- (367) Kuwabara K, et al. Relationship between Non-High-Density Lipoprotein Cholesterol and Low-Density Lipoprotein Cholesterol in the General Population. *J Atheroscler Thromb.* 2016; 23 (4): 477-90.
- (368) Nakamura M, Koyama I, Iso H, et al. Measurement performance of reagent manufacturers by Centers for Disease Control and Prevention/Cholesterol Reference Method Laboratory Network lipid standardization specified for metabolic syndrome-focused health checkups program in Japan. *J Atheroscler Thromb.* 16: 756-63. 2009.
- (369) Iso H, Naito Y, Sato S, et al. Serum triglycerides and risk of coronary heart disease among Japanese men and women. *Am J Epidemiol.* 153: 490-9. 2001.
- (370) 臨床検査のガイドラインJSLM2012. 日本臨床検査医学会. 2012.
- (371) 日本痛風・核酸代謝学会ガイドライン改訂委員会：高尿酸血症・痛風の治療ガイドライン改訂第3版編集. 2019.
- (372) Kanabrocki EL, Third JL, Ryan MD, et al. Circadian relationship of serum uric acid and nitric oxide. *JAMA.* 283: 2240-1. 2000.
- (373) Garrel DR, Verdy M, PetitClerc C, et al. Milk- and soy-protein ingestion: acute effect on serum uric acid concentration. *Am J Clin Nutr.* 53: 665-9. 1991.
- (374) Goldfarb DS, MacDonald PA, Hunt B, Gunawardhana L. Febuxostat in gout: serum urate response in uric acid overproducers and underexcretors. *J Rheumatol.* 2011; 38: 1385-9.
- (375) Law MR, et al. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ.* 2009; 338: b1665.
- (376) Trunbull F, et al. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-designed overviews of randomized trials. *Lancet.* 2003; 362: 1527-1535.
- (377) Staessen JA, et al. Cardiovascular prevention and blood pressure reduction: a quantitative overview updated until 1 March 2003. *J Hypertens.* 2003; 21: 1055-1076.
- (378) 日本高血圧学会高血圧治療ガイドライン作成委員会編. 高血圧治療ガイドライン2019. 日本高血圧学会, 2019.
- (379) Miura K, Nagai M, Ohkubo T. Epidemiology of hypertension in Japan: where are we now? *Circ J* 2013; 77: 2226-31.
- (380) Mori H, Ukai H, Yamamoto H et al. Current status of antihypertensive prescription and associated blood pressure control in Japan. *Hypertens Res* 2006; 29: 143-51.
- (381) Wald DS, et al. Combination therapy versus monotherapy in reducing blood pressure: meta-analysis on 11,000 participants from 42 trials. *Am J Med.* 2009; 122: 290-300.
- (382) 動脈硬化性疾患予防ガイドライン2017年版 第4章包括的リスク管理1動脈硬化性疾患の絶対リスクと脂質管理目標 日本動脈硬化学会(編)p.49-57 2017年 東京.

- (383) Nakamura H, Arakawa K, Itakura H, et al. for MEGA study group: Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA study): a prospective randomized controlled trial. Lancet. 368: 1155-63, 2006.
- (384) Scandinavian simvastatin survival study group: Randomized trial of cholesterol lowering in 4444 patients with coronary heart disease; the Scandinavian simvastatin survival study (4S). Lancet. 344: 1383-89, 1994.
- (385) 動脈硬化性疾患予防ガイドライン2017年版 第4章包括的リスク管理3薬物療法 日本動脈硬化学会(編) p.81-96 2017年 東京.
- (386) Ballantyne CM, Houri J, Notarbartolo A, et al.: the ezetimibe study group: Effect of ezetimibe coadministered with atorvastatin in 628 patients with primary hypercholesterolemia: a prospective, randomized, double-blind trial. Circulation, 107: 2409-15, 2003.
- (387) Kastelin JJ, Akdim F, Stroes ES, Zwinderman AH, et al. for the ENHANCE investigators: Simvastatin with or without ezetimibe in familial hypercholesterolemia. N Engl J Med. 358:1431-43, 2008.
- (388) Thomopoulos C, Skalis G, Michalopoulou H, et al. Effect of low-density lipoprotein cholesterol lowering by ezetimibe/simvastatin on outcome incidence: overview, meta-analysis, and meta-regression analyses of randomized trials. Clin Cardiol 2015; 38: 763-769.
- (389) Yokoyama M, Origasa H, Matsuzaki M, et al. for the Japan EPA lipid intervention study (JELIS) investigators: Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomized open-label, blinded endpoint analysis. Lancet. 369: 1090-9, 2007.
- (390) Ni YG, Condra JH, Orsatti L, et al. A proprotein convertase subtilisin-like / kexin type 9 (PCSK9) C-terminal domain antibody antigen-binding fragment inhibits PCSK9 internalization and restores low density lipoprotein up take. J Biol Chem 2010; 285: 12882-12891.
- (391) 日本痛風・核酸代謝学会ガイドライン改訂委員会：高尿酸血症・痛風の治療ガイドライン改訂第3版編集. 2019.

- (392) www.glycemicindex.com
- (393) Different trends in serum cholesterol levels among rural and urban populations aged 40-59 in Japan from 1960 to 1990. Okayama A, Ueshima H, Marmot MG, Elliott P, Yamakawa M, Kita Y. *J Clin Epidemiol.* 1995 Mar; 48 (3): 329-37.
- (394) Epidemiological studies in the Upernivik district, Greenland. Incidence of some chronic diseases 1950-1974. Kromann N, Green A. *Acta Med Scand.* 1980; 208 (5): 401-6.
- (395) 内閣府食品安全委員会. 平成18年度食品安全確保総合調査：「食品に含まれるトランス脂肪酸の評価基礎資料調査報告書」.
- (396) The effects of different food fats on serum cholesterol concentration in man. ANDERSON JT, KEYS A, GRANDE F. *J Nutr.* 1957 Jul 10; 62 (3): 421-4. No abstract available.
- (397) Dietary protein sources and the risk of stroke in men and women. Bernstein AM, Pan A, Rexrode KM, Stampfer M, Hu FB, Mozaffarian D, Willett WC.
- (398) Panel on Dietary Reference Intakes for Electrolytes and Water SCotSEoDRI. Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate. Washington, DC: The National Academies Press 2005.
- (399) 奥田奈賀子・岡山明. ナトリウム・カリウム摂取と循環器疾患. *循環器内科* 2011; 70 (6): 546-54.
- (400) WHO. Guideline: Sodium intake for adults and children 2012 [Available from: http://www.who.int/nutrition/publications/guidelines/sodium_intake/en/index.html accessed March 9 2018.]
- (401) 厚生労働省. 日本人の食事摂取基準2015年 2014 [Available from: <http://www.mhlw.go.jp/stf/shingi/0000041824.html> accessed March 12 2018.]
- (402) 日本高血圧学会. 高血圧治療ガイドライン 2014 [Available from: <https://www.jpnsh.jp/guideline.html> accessed March 13 2015.]
- (403) 厚生労働省. 平成28年国民健康・栄養調査報告 2017 [Available from: <http://www.mhlw.go.jp/bunya/kenkou/eiyou/h28-houkoku.html> accessed March 9 2018.]
- (404) Krauss RM, Eckel RH, Howard B, et al. AHA Dietary Guidelines: revision 2000: A statement for healthcare professionals from the Nutrition Committee of the American Heart Association. *Circulation* 2000; 102 (18): 2284-99. [published Online First: 2000/11/01]
- (405) Young DB, Lin H, McCabe RD. Potassium's cardiovascular protective mechanisms. *The American journal of physiology* 1995; 268 (4 Pt 2): R825-37. [published Online First: 1995/04/01]
- (406) He FJ, MacGregor GA. Beneficial effects of potassium on human health. *Physiologia plantarum* 2008; 133 (4): 725-35. [published Online First: 2008/08/30]
- (407) 厚生労働省. 平成28年国民健康・栄養調査報告 2017 [Available from: <http://www.mhlw.go.jp/bunya/kenkou/eiyou/h28-houkoku.html> accessed March 9 2018.]
- (408) Panel on Dietary Reference Intakes for Electrolytes and Water SCotSEoDRI. Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate. Washington, DC: The National Academies Press 2005.
- (409) 日本腎臓学会. 慢性腎臓病に対する食事療法基準2014年版. 東京: 東京医学社 2014.
- (410) WHO. Guideline: Potassium intake for adults and children 2012 [Available from: http://www.who.int/mediacentre/news/notes/2013/salt_potassium_20130131/en/ accessed July 31th 2013.]
- (411) 厚生労働省. 日本人の食事摂取基準2015年 2014 [Available from: <http://www.mhlw.go.jp/stf/shingi/0000041824.html> accessed April 9 2016.]

- (412) 文部科学省科学技術・学術審議会資源調査分科会. 日本食品標準成分表2015年版(七訂). 東京: 全国官報販売協同組合 2015.
- (413) Okuda N, Okayama A, Miura K, et al. Food sources of dietary sodium in the Japanese adult population: the international study of macro-/micronutrients and blood pressure (INTERMAP). European journal of nutrition 2017; 56 (3): 1269-80. doi: 10.1007/s00394-016-1177-1.
- (414) 厚生労働省. 日本人の食事摂取基準2015年 2014 [Available from: <http://www.mhlw.go.jp/stf/shingi/0000041824.html> accessed March 12 2018.]
- (415) WHO. Guideline: Potassium intake for adults and children 2012 [Available from: http://www.who.int/nutrition/publications/guidelines/potassium_intake/en/ accessed]
- (416) 日本病院薬剤師会雑誌, 第40巻8号, P951-953, 2004.
- (417) Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. Omenn GS, Goodman GE, Thorquist MD, Balmes J, Cullen MR, Glass A, Keogh JP, Meyskens FL Jr, Valanis B, Williams JH Jr, Barnhart S, Cherniack MG, Brodkin CA, Hammar S.J Natl Cancer Inst. 1996 Nov 6; 88 (21): 1550-9.
- (418) Appel LJ, Champagne CM, Harsha DW, Cooper LS, Obarzanek E, Elmer PJ, Stevens VJ, Vollmer WM, Lin PH, Svetkey LP, Stedman SW, Young DR; Writing Group of the PREMIER Collaborative Research Group. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. JAMA. 2003; 289: 2083-93.
- (419) 板井一好, 奥田奈賀子, 岡山明. 岩手県内における主な外食に含まれる食塩量について. 第52回日本循環器病予防学会学術集会; 2016 2016年6月; さいたま市.
- (420) Okuda N, Okayama A, Miura K, et al. Food sources of dietary sodium in the Japanese adult population: the international study of macro-/micronutrients and blood pressure (INTERMAP). European journal of nutrition 2017; 56 (3): 1269-80. doi: 10.1007/s00394-016-1177-1.
- (421) 適塩・血圧対策推進協会. ナト・カリ塩とは [Available from: <http://www.lowsalt.or.jp/natokari.html> accessed January 12 2018.]
- (422) 日本体力医学会体力科学編集委員会・監訳:「1. 身体活動に伴う有益性とリスク」, 運動処方の指針－運動負荷試験と運動プログラム－, 原書第8版, p8-9, 2011, 南江堂, 東京.
- (423) 石井直方:スロートレーニング(特集 常識を打ち破る運動生理学の新知見). 体育の科学, 59(3), 177-183, 2009.
- (424) 河村顕治:「24変形膝関節症」, 運動療法ガイドー正しい運動処方を求めて, 井上一, 武藤芳照, 福田潤編著, 改訂第4版, p348-356, 2006, 南江堂, 東京.
- (425) Hambrecht RP, Schuler GC, Muth T: Greater diagnostic sensitivity of treadmill versus cycle exercise testing of asymptomatic men with coronary artery disease. Am J Cardiol, 70: 141-146, 1992.
- (426) Myers J, Buchanan N, Walsh D: Comparison of the ramp versus standard exercise protocols. J Am Coll Cardiol, 17: 1334-1342, 1991.
- (427) Gellish RL, Goslin BR, Olson RE, McDonald A, Russi GD, Moudgil VK: Longitudinal modeling of the relationship between age and maximal heart rate. Med Sci Sport Exer, 39: 822-829, 2007.
- (428) 村瀬訓生, 勝村俊仁, 上田千穂子, 井上茂, 下光輝一:身体活動量の国際標準化—IPAQ日本語版の信頼性, 妥当性の評価—, 厚生の指標, 49(11): 1-9, 2002.
- (429) Craig C.L., Marshall A.L., Sjöström M., Bauman A.E., Booth M.L., Anisworth B.E., et al.

- International physical activity questionnaire: 12-country erliability and validity. Med Sci Sports Exerc. 35, 1381-1395. 2003.
- (430) 池上晴夫：運動処方－理論と実際－，新版，p147，1990，朝倉書店，東京。
- (431) 運動基準・運動指針の改定に関する検討会：「健康づくりのための身体活動基準2013」，運動基準・運動指針の改定に関する検討会報告書，2013.<<http://www.mhlw.go.jp/stf/houdou/2r9852000002xpxpqt.pdf>>
- (432) 厚生労働省：「日本人の食事摂取基準(2015年版)」策定検討会報告書。<<http://www.mhlw.go.jp/file/05-Shingikai-10901000-Kenkoukyoku->>
- (433) Ishikawa-Takata K, Tabata I, Sasaki S, et al. Physical activity level in healthy free-living Japanese estimated by doubly labelled water method and International Physical Activity Questionnaire. Eur J Clin Nutr, 62:885-891, 2008.
- (434) Ishikawa-Takata K, Naito Y, Tanaka S, et al. Use of doubly labeled water to validate a physical activity questionnaire developed for the Japanese population. J Epidemiol, 21: 114-121, 2011.
- (435) Tambalis K, et al.: Responses of blood lipids to aerobic, resistance, and combined aerobic with resistance exercise training : a systematic review of current evidence. Angiology, 60: 614-632, 2009.
- (436) Miyachi M, et al.: Unfavorable effects of resistance training on central arterial compliance: a randomized intervention study. Circulation, 110: 2858-2863, 2004.
- (437) Okamoto T, et al.: Effects of eccentric and concentric resistance training on arterial stiffness. J Hum Hypertens, 20: 348-354, 2006.
- (438) Yamamoto K, et al.: Poor trunk flexibility is associated with arterial stiffening. Am J Physiol Heart Circ Physiol, 297: H1314-H1318, 2009.
- (439) 野村武雄, 金田晃一：i. アクアエクササイズ, 高齢社会における運動支援実践ガイド. 臨床スポーツ医学, 22(臨時増刊号) : 150-157, 2005.
- (440) 柳田昌彦, 交野好子：福井県における楽しい集団的介護予防体操「ふくいイッショライダンベル体操」の創作. 北陸公衆衛生学会誌, 33(2) : 60-69. 2007.
- (441) 柳田昌彦, 塩津陽子, 炭本佑佳：5年間にわたる「ふくいイッショライダンベル体操」の定期的実践が地域在住高齢女性の生活体力, 形態および健康関連QOLに及ぼす効果—福井県敦賀市の介護予防地域支援事業に介入して—. ライフケアジャーナル, 7(1) : 7-14, 2016.
- (442) Br Med J. 1964 Jun 6; 1 (5396): 1460-7 MORTALITY IN RELATION TO SMOKING: TEN YEARS' OBSERVATIONS OF BRITISH DOCTORS. DOLL R, HILL AB.
- (443) 生活習慣病予防研究センター. 簡易版食傾向調査票, 生活調査票「知食スマート版®」2019 [Available from: <http://www.jrisp.com/shg/>] accessed August 30 2019.]
- (444) Okuda N, Itai K, Okayama A. Usefulness of a short dietary propensity questionnaire in Japan. J Atheroscler Thromb (accepted on Dec 11, 2017) 2017.
- (445) Prochaska JO, et al. The transtheoretical model of health behavior change. Am J Health Promot 1997; 38-48.
- (446) Prochaska JO, et al. "The transtheoretical model and stages of change", Health behavior and health education: theory, research, and practice. 4th ed. Glanz K, et al. eds. Jossey-Bass 2008; 97-121.
- (447) 岡山明編著, 保健同人社. 生活習慣改善シリーズ メタボリックシンドローム予防の健康教育教材を用いた実践的プログラム. 2007; 175.
- (448) 個情第752号, 基発0529第6号平成29年5月29日 雇用管理分野における個人情報のうち健康情報を取り扱うに当たっての留意事項について.